

Improving Feed Quality, Availability, and Cost with Alternative Feeds and Feeding Regimes for Smallholder Aquaculture Operations

Morgan Chow, Briana Goodwin,
Stephanie Ichien, Kat Goetting,
and *Hillary Egna



Feed the Future Innovation Lab for
Collaborative Research on Aquaculture & Fisheries
Oregon State University, Corvallis, OR 97331
AquaFish@oregonstate.edu
Website: aquafish.oregonstate.edu

Introduction

Aquaculture growth for smallholder farms is often limited by a lack of access to affordable, high-quality feed sources. Furthermore, feed costs constitute the majority of production costs which are estimated to be between 60-80% for tilapia, one of the most widely cultured species in the world. Research supported by the AquaFish Innovation Lab in Africa and Asia is **focusing on reducing operational costs and negative environmental impacts by developing alternative feed types and feeding regimes** while maintaining high yields.




More specifically, research involves investigating affordable and locally sourced feeds and **examining how pulsed feeding strategies affect fish gut biomes** to improve nutrient uptake efficiency from feeds. In doing so, AquaFish researchers are improving feed quality and availability and reducing reliance on expensive feeds, which will increase productivity and sustainability of small-scale aquaculture operations in communities across the globe.

Feed Ingredients

AquaFish researchers have tested alternative feed ingredients that are locally sourced and more affordable than their expensive resource intensive fishmeal counterparts.



At **Sokoine University of Agriculture in Morogoro, Tanzania**, AquaFish researchers investigated invertebrates – housefly larvae and earthworms – for use in fish feeds as they both have short life-cycles and high fecundity. Results produced a cost-effective and protein-efficient composition for tilapia production.



The chemical composition of maggots (*Musca domestica*), earthworms (*Lumbricus terrestris*), and fish meal were determined for crude protein, fat, crude fiber, and ash. Based on results, nine feed formulations were developed using mixtures of fish meal, maggot meal (HM), earthworm meal (EWM), and cotton seed cake. A feeding experiment was conducted to evaluate the growth performance, feed utilization, and cost effectiveness of the nine feeds on Nile tilapia. Growth, feed utilization, and cost was most optimal in fish fed diets with 35% HM and EWM and 5% fish meal or plant protein.



At **Can Tho University, in Vietnam** and the **University of Rhode Island**, AquaFish researchers developed alternative feeds for small-scale snakehead farming with soybean, rice bran, and cassava, to reduce fishmeal content in commercial diets.

Feed Strategies

Groundbreaking AquaFish feeds research at **Central Luzon State University** in the **Philippines** found that feeding tilapia on alternate days was effective at producing tilapia at similar yields as if fed at a full daily ration.

These findings have served as the baseline for other AquaFish studies that have assessed the outcomes of alternate-day feeding for other fish species as well as polyculture systems. If alternate-day feeding can produce equivalent fish production rates, while also cutting feed costs in half and reducing nutrient inputs to the environment, then it is a worthwhile strategy to promote for aquaculture.



Impacts on nutrient availability and beneficial gut flora of fish

Background



- Improving the understanding of how finfish acquire and utilize nutrient inputs is essential to future improvements in aquaculture production efficiency.
- Feed ingredients and feeding strategies can influence the diversity and abundance of intestinal microbiota in both humans and fish.
- Reduced feeding can potentially enhance nutrient uptake efficiency or promote foraging on in-pond primary production, diversifying the fish's diet and enhancing nutrient recycling in ponds.

Research in the Philippines demonstrated that equivalent production yields can be achieved with much less feed (50% reduction) through the implementation of **pulsed feeding strategies**. Follow-up experiments at **Bangladesh Agricultural University** and **North Carolina State University** examined how alternate-day feeding strategies may enhance nutrient absorption by measuring nutrient transporter abundance and gut microbial diversity in response to different feeding regimes.

Feed variations were tested according to their impact on the:

- Growth performance,
- Gastrointestinal nutrient absorption efficiency, and
- Establishment of beneficial gut flora for tilapia pond culture.



Results

Metagenomic analyses identified 145 different families of prokaryotic (all bacteria) and 132 eukaryotic organisms in the fecal material of tilapia (Figures 1 and 2). The highest diversity of eukaryotes was found in **fish fed every other day with pond fertilization**.

There were no significant differences in gene expression of these nutrient transporters in any treatment that incorporated both feeding and pond fertilization. Expression of the transporters was higher in fish that were not fed and grown in fertilized ponds and lower in fish that were fed daily without pond fertilization. Fish fed on alternate days had more moderate expression levels of certain transporters which may allow for more balanced and efficient nutrient uptake.

Together, these studies along with those in the Philippines indicate **feeding Nile tilapia on alternate days along with weekly pond fertilization has no negative effects on growth, survivability, or production** compared to daily feeding regimes, and additionally produces the greatest net return on investments.

This work also suggests for the first time that **combined feeding and fertilization produces the greatest biodiversity of microbiomes in the intestine** which could contribute to enhanced feed efficiency and overall health of tilapia, particularly those subjected to more moderate feeding strategies. This work lays the framework for development of probiotic supplements that can be incorporated into diet formulations for improving growth and nutrient absorption.

Treatment / Factors	Daily feeding with 4:1 (N:P) weekly	Feeding every other day 4:1 (N:P) weekly	Feeding every 3rd day 4:1 (N:P) weekly	No feeding 4:1 (N:P) weekly	Daily feeding No fertilization
Stocking Density	5 fish/ m ²	5 fish/ m ²	5 fish/ m ²	5 fish/ m ²	5 fish/ m ²
Initial Weight (g)	3.55±0.90 ^a	3.55±0.90 ^a	3.55±0.90 ^a	3.55±0.90 ^a	3.55±0.90 ^a
Final Weight (g)	127.63±2.75 ^a	120.17±5.44 ^a	85.10±11.13 ^b	43.15±4.28 ^c	129.53±8.59 ^a
Weight Gain (g)	124.08±2.75 ^a	116.62±5.44 ^a	81.55±11.13 ^b	39.60±4.28 ^c	125.98±8.59 ^a
SGR (%/day)	3.14±0.02 ^a	3.09±0.04 ^a	2.78±0.12 ^b	2.19±0.09 ^c	3.15±0.06 ^a
FCR	1.44±0.10 ^a	0.93±0.09 ^b	0.68±0.15 ^c	—	1.61±0.10 ^c
Survival Rate (%)	93.44±6.26 ^a	91.66±8.00 ^a	90.70±9.74 ^{ab}	76.79±2.68 ^b	97.71±2.11 ^a
Production (kg/pond)	92.98±10.29 ^{ab}	86.40±19.33 ^a	58.93±7.17 ^c	27.50±4.57 ^d	102.63±10.66 ^a
Net Production (kg/ha)	4091.42±354.52 ^a	5658.26±527.83 ^a	4086.39±640.07 ^a	1796.04±233.89 ^b	4392.81±461.41 ^a
Total Production (kg/ha)	6292.09±354.52 ^a	5837.84±527.83 ^a	4179.08±640.07 ^a	1950.35±233.89 ^b	6578.53±461.41 ^a

Table 3: Growth factors according to five different treatments. Values with different letters are significantly different (P<0.05).¹




Figure 1: Relative abundance of higher groups of eukaryotes in the fecal material of Nile tilapia subjected to varying pulsed-feeding regimes.^{1,2,3}




Figure 2: Relative abundance of bacteria in the fecal material of Nile tilapia subjected to varying pulsed-feeding regimes at the level of Family/Species.^{1,2,3}




Figure 3: Predicted metabolic function of gut flora that may affect metabolism and nutrition in fish.³

AquaFish is grateful to participants involved in feed ingredients and feed strategies work in Bangladesh, Philippines, Tanzania, and Vietnam, in particular for their field research and analysis -- much of which is available in the peer-reviewed literature below:

Be, T.T. and T.T.T. Hien. 2010. Replacement of fish meal protein by soybean meal protein with or without phytase supplementation in snakehead (*Channa striata*) diets. The Scientific Journal of Can Tho University.

Bolivar, R.B., E.B.T. Jimenez, R.M.V. Sayco, R.L.B. Argueza, C.R. Stark, and P.R. Ferket. 2011. Fishmeal-Free Diets Improve the Cost Effectiveness of Culturing Nile Tilapia (*Oreochromis niloticus* L.) in Ponds Under an Alternate Day Feeding Strategy. Better Science, Better Fish, Better Life: Proceedings of the Ninth International Symposium on Tilapia in Aquaculture (2011) [Edited By: Liu Liping and Kevin Fitzsimmons] pg: 268-274.

Borski, R.J., R.B. Bolivar, E.B.T. Jimenez, R.M.V. Sayco, R.L.B. Argueza, C.R. Stark, and P.R. Ferket. 2011. Fishmeal-Free Diets Improve the Cost Effectiveness of Culturing Nile Tilapia (*Oreochromis niloticus* L.) in Ponds Under an Alternate Day Feeding Strategy. Better Science, Better Fish, Better Life: Proceedings of the Ninth International Symposium on Tilapia in Aquaculture (2011) [Edited By: Liu Liping and Kevin Fitzsimmons] pg: 95-101.

Gurung, S., M.K. Shrestha, and N.P. Pandit. 2013. Our Nature 11(2): 116-125.

Grim-Greenblatt, J., R. Pomeroy, B. Bravo-Ureta, L.X. Sinh, H.V. Hien, and T. Getchis. 2015. Economic analysis of alternative snakehead *Channa striata*

feed. Aquaculture Economics & Management 19(2): 192-209.

Hien, T.T.T., T.T. Be, C.M. Lee, and D.A. Bengtson. 2015. Development of formulated diets for snakehead (*Channa striata* and *Channa micropeltes*): Can phytase and taurine supplementation increase use of soybean meal to replace fish meal? Aquaculture 448: 334-340.

Hien, T.T.T., L.Q. Toan, T.T. Be, and N.H.D. Trung. 2010. Replacing fish meal by soybean meal in giant snakehead (*Channa micropeltes*) diets. The Scientific Journal of Can Tho University.

Hien, T.T.T., T.L.C. Tu, N.V. Tien, N.B. Trung, T.M. Phu, P.M. Duc, and D. Bengtson. 2014. Replacing fishmeal by some of soy protein sources in feed for snakehead (*Channa striata*). Can Tho University Journal of Science 1: 310-318.

Kasiga, T., R. Chen, T. Sink, and R. Lochmann. 2014. Effects of Reduced Soybean-Meal Diets Containing Moringa oleifera or Leucaena leucocephala Leaf Meals on Growth Performance, Plasma Lysozyme, and Total Intestinal Proteolytic Enzyme Activity of Juvenile Nile Tilapia, *Oreochromis niloticus*, in Outdoor Tanks. Fisheries Science 45(5): 508-522.

Nam, S. 2009. Snakehead aquaculture in the Mekong Delta, Vietnam. Cambodia Fisheries Magazine 13.

Shrestha, M.K., R.L. Sharma, K. Gharti, and J.S. Diana. 2011. Aquaculture 319: 284-289.

Singh, L.X. and R.S. Pomeroy. 2010. Current situation and challenges for farming of snakehead fish (*Channa micropeltes* and *Channa striatus*) in the Mekong Delta, Vietnam. Sustainable Aquaculture 15(4): 11-17.

Vera Cruz, E.M. 2009. Influence of the photoperiod on growth rate and insulin-like growth factor-I gene expression in Nile Tilapia *Oreochromis niloticus*. Journal of Fish Biology 75: 130-141.

Yi, Y., and J. Diana. 2008. Strategies for Nile Tilapia (*Oreochromis niloticus*) Pond Culture. Proceedings of the 8th International symposium on Tilapia in Aquaculture, Cairo, Egypt. 12-14 October 2008

1. Salger, S., Reza, J. Baltzegar, A., Haque, S. M., Wahab, M. A., and Borski, R. (2016). Assessment of pulsed feeding strategies on growth performance, gastrointestinal nutrient absorption efficiency, and establishment of beneficial gut flora for tilapia pond culture. In AquaFish Innovation Lab (ed). AquaFish CRSP Technical Reports: Investigations 2013-2015. Corvallis, OR.

2. Borski, R. 2016. Characterization of gut microbial communities associated with improved feed efficiency in tilapia pond culture. [Oral presentation]. 2016 AquaFish Program Meeting, Surabaya, Indonesia. 24-25 April 2016.

3. Borski, R., S. Salger, D. Baltzegar, J. Reza, and Md. Abdul Wahab. 2016. Effect of pulsed feeding on growth, gut metagenome, and intestinal nutrient transporters of tilapia in pond culture. [Oral presentation]. 2016 Asia-Pacific Aquaculture, Surabaya, Indonesia, 27 April 2016.

Citations:

