Notice of Publication

AQUACULTURE COLLABORATIVE RESEARCH SUPPORT PROGRAM

RESEARCH REPORTS

Sustainable Aquaculture for a Secure Future

Title:

Comparative and evolutionary analysis in natural diploid and tetraploid weather loach Misgurnus anguillicaudatus based on cytochrome b sequence data in central China

Author(s):

Chengtai Yang¹, Ling Cao1, Weimin Wang¹ Contact Information, Yi Yang², Khalid Abbas¹, Binlun Yan³, Hanping Wang⁴, Li Su⁵, Yuhua Sun¹ and Huanling Wang¹

¹College of Fisheries, Key Lab of Agricultural Animal Genetics, Breeding and Reproduction of Ministry of Education, Huazhong Agricultural University, Wuhan, Hubei, 430070, People's Republic of China

²Aquaculture and Aquatic Resources Management, School of Environment, Resources and Development, Asian Institute of Technology, P.O. Box 4, Klong Luang, Pathum Thani, 12120, Thailand

³Jiangsu Key Lab of Marine Biotechnology, Huaihai Institute of Technology, Lianyungang, Jiangsu, 222005, People's Republic of China

⁴Aquaculture Genetics and Breeding Laboratory, Ohio State University South Centers, 1864 Shyville Road, Piketon, OH 45661, USA

⁵Huangshi Institute of Technology, Huangshi, Hubei, 435003, People's Republic of China

Date: August 18, 2008

Publication Number: CRSP Research Report 07-227

The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.

Abstract:

To obtain the phylogenetic relationship between diploid and tetraploid $Misgurnus\ anguillicaudatus$, the mitochondrial cyt b gene in the diploid and tetraploid weather loach were isolated and sequenced. The DNA sequences were analyzed using MEGA 3.0 software to determine the phylogenetic relationship. Forty-five variable sites among cyt b gene sequences and 18 amino acid substitutions occurred within the diploid and tetraploid loaches as deduced from the nucleotide sequences analysis of the cyt b gene. The nucleotide pairwise distance between diploid and tetraploid loach ranged from 0.001 to 0.025. Phylogenetic analysis revealed evolutionary relationships between diploid and tetraploid loach. Our results indicated a significant difference between diploid and tetraploid loach about the cyt b gene. AMOVA analysis indicated that there were no significant genetic variations within diploid loaches (Fst = 0.2529, P>0.05) and within tetraploid loaches (Fst = 0.0564, P<0.05), neither. However, significant genetic differences were found between diploid and tetraploid loaches (Fst = 0.7634, P<0.05). Thus, it is concluded that no reproductive isolation was found within the same cytotypes of different localities, but there was reproductive isolation between these two cytotypes. The diploid loach existed before the tetraploid loaches in nature. The pres-

Continued...

ent study is first to describe the phylogenetic relationships of nature polyploidy weather loach using mtDNA cyt b gene.

This abstract is excerpted from the original paper, which was published online in Environmental Biology of Fishes DOI: 10.1007/s10641-007-9283-9, 09/20/07.