

NOTICE OF PUBLICATION

RESEARCH REPORTS

TITLE XI POND DYNAMICS/AQUACULTURE COLLABORATIVE RESEARCH SUPPORT PROGRAM

- Title:** An analysis of biological characteristics of *Macrobrachium rosenbergii* (de Man) in relation to pond production and marketing in Thailand
- Author(s):** C. Kwei Lin, Agricultural and Food Engineering Division, Asian Institute of Technology, P.O. Box 2754, Bangkok, Thailand; Great Lakes Research Division, The University of Michigan, Ann Arbor, Michigan 48109, USA
Mali Boonyaratpalin, National Inland Fisheries Institute, Bangkok, Thailand
- Date:** 14 February 1989 **Publication Number:** CRSP Research Reports 89-16
- Price:** The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.
- Abstract:** Data on production and marketing of giant freshwater prawn were collected over a seven-month, grow-out period through collaboration with a medium-sized commercial prawn farm in central Thailand. Juvenile prawns with an average weight of 4.2 g were stocked at a density of 6 prawns/m² in three 0.5-ha earthen ponds. Average growth rate determined during the first three months of the grow-out period was 0.4 g/prawn per day; prawns of marketable size were harvested selectively during the remaining four months of the rearing period, resulting in a total accumulated yield of 1.3 tons/ha with an average prawn weight of 32 g and 60% survival. As different sexes and sizes of prawns were sold at different prices, the harvests were customarily sorted into several categories: large, medium, and small males, long-clawed males, soft shells, females with eggs, females without eggs, and terminal males. The total weight and number of prawns recorded for each of these categories showed that the female to male ratios were 1.6:1 and 4:1 by weight and number, respectively, while the ratio of short- to long-clawed males was 3:1 by weight and 4:1 by number. Four percent of the marketable population was termed "soft shells" and 64% of the females bore eggs. The ratio of head weight to tail weight of marketable prawns varied substantially among the different categories: 1.0:1 for females, 2.5:1 for long-clawed males, and 1.6:1 for short-clawed males. Females predominated in the first and second 1.5-month harvest periods while males predominated in the final 1.5 months of the harvest. The economic yield of prawn culture was not only determined by the biomass, but also by the population structure of the various biological categories and the harvest season.

This paper was published in *Aquaculture*, 74 (1988) 205-215.

CRSP RESEARCH REPORTS are published as occasional papers by the Program Management Office, Pond Dynamics/Aquaculture Collaborative Research Support Program, Office of International Research and Development, Oregon State University, Snell Hall 400, Corvallis, Oregon 97331-1641 USA. The Pond Dynamics/Aquaculture CRSP is supported by the U.S. Agency for International Development under CRSP Grant No.: DAN-4023-G-SS-7066-00.