

NOTICE OF PUBLICATION



Title: Nitrogen input, primary productivity and fish yield in fertilized freshwater ponds in Indonesia

Author(s): Christopher F. Knud-Hansen, Ted R. Batterson, Clarence D. McNabb
Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI 48824, USA

Irwan S. Harahat, Komar Sumantadinata and H. Muhammed Eldman
Institut Pertanian Bogor, Fakultas Perikanan, Jalan Raya Pajajaran, Bogor, Indonesia

Date: 27 July 1991 **Publication Number:** CRSP Research Reports 91-32

Price: The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.

Abstract: Twelve 0.02-ha ponds in West Java were fertilized weekly with four levels of chicken manure (12.5, 25, 50, and 100 g dry weight/m²/week) during a 149-day growout experiment for Nile tilapia (*Oreochromis niloticus*) production. Laboratory leaching experiments for measuring dissolved inorganic nitrogen (DIN) release from chicken manure showed that nitrogen was released as ammonia-N, which was rapidly lost from the manure and leveled off at about 6 mg NH₄-N/g dry weight chicken manure after 4-5 days. Allochthonous DIN input from both chicken manure fertilization and almost daily source-water additions ranged from 0.055 to 0.142 g N/m²/day. Source water contributed more nitrogen than manure in all but the highest fertilization treatment.

Pond averages of net primary productivity (NP) ranged from 0.54 to 2.00 g C/m²/day, while gross fish yield at harvest ranged from 4.9 to 15.7 kg fresh weight/ha/day. Net fish yield (NFY) was linearly correlated to both the dry weight sum of NP and chicken manure fertilization ($r^2=0.97$) and allochthonous DIN input ($r^2=0.96$). Results suggest that Nile tilapia obtained organic carbon from both primary productivity and manure-derived detritus.

DIN availability limited algal productivity at a chicken manure fertilization rate of 100 g dry weight/m²/week when microbial decomposition of manure supplied sufficient CO₂. Incorporation of allochthonous DIN input into NFY increased significantly from 15.0% at the three lower fertilization rates to 25.4% at the highest loading rate. Using organic fertilizers proportionally rich in phosphorus and carbon relative to nitrogen may maintain this element's limitation of algal productivity. Efficient utilization of DIN input also may minimize total and un-ionized ammonia concentrations. In the absence of deleterious ammonia effects on survival and growth, fish yields can be readily predicted from measurements of nitrogen inputs.

This abstract was reprinted from the original, which was published in *Aquaculture*, 94 (1991)49-63, Elsevier Science Publishers B.V., Amsterdam

CRSP RESEARCH REPORTS are published as occasional papers by the Program Management Office, Pond Dynamics/Aquaculture Collaborative Research Support Program, Office of International Research and Development, Oregon State University, Snell Hall 400, Corvallis, Oregon 97331-1641 USA. The Pond Dynamics/Aquaculture CRSP is supported by the U.S. Agency for International Development under CRSP Grant No.: DAN-4023-G-00-0031-00.